

Presented at the
24th Annual Conference on
Physical Electronics
M.I.T., Cambridge, Mass.,
March 25-27, 1964

CPST
HC. 1.00
ME. .50

FACILITY FORM 6

(ACCESSION NUMBER)	(THRU)
10	1
(PAGES)	(CODE)
TMX-51612	26
(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

AN ANALYTICAL STUDY OF THE WORK FUNCTION CHARACTERISTICS OF A METAL
IMMERSED IN CESIUM VAPOR - SURFACE CONTRIBUTION

By Keung P. Luke

Lewis Research Center

National Aeronautics and Space Administration

Cleveland, Ohio, 44135

It is well known that adsorbed cesium significantly reduces the electron work function of the emitter and collector in a cesium thermionic diode.

Figure 1, which is based on the data reported by Houston¹ of General Electric, shows how this reduction in electron work function, $-\Delta\phi_e$, varies with the ratio of metal surface temperature to cesium reservoir temperature, T_s/T_{cs} .

In an effort to derive an analytic expression to represent data of the type shown in Fig. 1, it is found that the metal surface plays a far more important role in the adsorption process than that attributed to it in the literature. It will be shown that because the presence of the metal surface, it is necessary to consider the effective electric field acting on each adsorbed particle, the effective polarizability of each adsorbed particle, and a new expression for the variation in atom desorption energy with gas coverage.

Consider the model of an adsorbed particle shown in Fig. 2. Let $(\vec{E}_{eff})_i$ be the effective dipole field acting on the i th adsorbed particle. Because of this field, the effective dipole moment $(\vec{m}_{eff})_i$ of this particle is changed by $2\alpha_i(\vec{E}_{eff})_i$ to yield a new effective dipole moment $(\vec{p}_{eff})_i$ given by the relation

$$(\vec{p}_{eff})_i = (\vec{m}_{eff})_i + 2\alpha_i(\vec{E}_{eff})_i \quad (1)$$

The effective dipole field $(\vec{E}_{eff})_i$ consists of the field \vec{E}_{ii} produced by the image of the dipole $\alpha_i(\vec{E}_{eff})_i$ induced in the i th adsorbed particle

Approved for release by NSA on 09-10-2013 pursuant to E.O. 13526

by $(\vec{E}_{\text{eff}})_i$, and the field $\sum_{\substack{k=1 \\ k \neq i}}^{\infty} \vec{E}_{ik}$ produced by all the other effective dipoles

(adsorbed particles), that is,

$$(\vec{E}_{\text{eff}})_i = \vec{E}_{ii} + \sum_{\substack{k=1 \\ k \neq i}}^{\infty} \vec{E}_{ik} \quad (2)$$

Assuming that the dipole field acting on each adsorbed particle is uniform and ^{is} equal to the value at the surface yields

$$\vec{E}_{ii} = 2 \frac{d_i (\vec{E}_{\text{eff}})_i}{(z_o)_i^3} \quad (3)$$

$$\vec{E}_{ik} = - \frac{(\vec{p}_{\text{eff}})_k}{r_{ik}^3} \quad (4)$$

Substituting equations (3) and (4) into equation (2) leads to the important new result

$$(\vec{E}_{\text{eff}})_i = - \frac{1}{1 - \frac{2 d_i}{(z_o)_i^3}} \sum_{\substack{k=1 \\ k \neq i}}^{\infty} \frac{(\vec{p}_{\text{eff}})_k}{r_{ik}^3} \quad (5)$$

Using the above expression for $(\vec{E}_{\text{eff}})_i$ in equation (1) yields the following equation for $(\vec{p}_{\text{eff}})_i$

$$(\vec{p}_{\text{eff}})_i = (\vec{m}_{\text{eff}})_i - \frac{2 d_i}{\left(1 - \frac{2 d_i}{(z_o)_i^3}\right)} \sum_{\substack{k=1 \\ k \neq i}}^{\infty} \frac{(\vec{p}_{\text{eff}})_k}{r_{ik}^3} \quad (6)$$

If the configuration of the effective dipoles associated with the adsorbed particles is assumed to be a Topping square array, equation (6) may be written as

~~THESE RESULTS ARE NOT VALID FOR A Topping SQUARE ARRAY~~

$$p_{\text{eff}} = \frac{\vec{m}_{\text{eff}}}{1 + 9.033 \mathcal{L}_{\text{eff}} (N_s \theta)^{3/2}} \quad (7)$$

where all dipoles are considered as identical and the effective polarizability \mathcal{L}_{eff} is defined as

$$\mathcal{L}_{\text{eff}} = \frac{2\mathcal{L}}{1 - \frac{2\mathcal{L}}{z_o^3}} \quad (8)$$

It should be noted that the new and unique feature in equation (7) is the use of the effective polarizability \mathcal{L}_{eff} in the denominator. For the cesium-tungsten system, analysis of Taylor and Langmuir's data² yields the value of 18.7 \AA^3 for the effective polarizability \mathcal{L}_{eff} of adsorbed cesium.

The atom desorption energy φ_a is the work $W_{s-\infty}$, needed to take an adsorbed particle from the surface to infinity in the form of an atom. Thus, it may be written

$$\varphi_a = W_{s-\infty} \quad (9)$$

or

$$\begin{aligned} \Delta \varphi_a &= \Delta W_{s-\infty} = - \Delta W_{\infty-s} \\ &= - \left[\frac{1}{2} \mathcal{L} |\vec{E}_{\text{eff}}|^2 - \left(\frac{p_{\text{eff}}}{2} \right) \cdot \vec{E}_{\text{eff}} \right] \end{aligned} \quad (10)$$

The first term within the bracket in equation (10) is the polarization energy, the second term is the adsorbed particles-effective dipole field interaction energy, and \vec{E}_{eff} is the effective dipole field given by equation (5). In terms of $|\vec{m}_{\text{eff}}|$, $|\vec{p}_{\text{eff}}|$, \mathcal{L}_{eff} and z_o equation (10) becomes

$$\Delta \varphi_a = - \frac{1}{4 \mathcal{L}_{\text{eff}}} \left(1 + \frac{\mathcal{L}_{\text{eff}}}{z_o^3} \right) \left(|\vec{m}_{\text{eff}}|^2 - |\vec{p}_{\text{eff}}|^2 \right) \quad (11)$$

Since

$$\Delta\psi_e = - 2 \pi e N_s (\vec{p}_{\text{eff}} \cdot \hat{n}) \theta \quad (12)$$

and

$$\vec{p}_{\text{eff}} \cdot \hat{n} = \pm |\vec{p}_{\text{eff}}| \quad (13)$$

equation (11) may be written in the form

$$\Delta\psi_a = - \frac{1}{4 \mathcal{L}_{\text{eff}}} \left(1 + \frac{\mathcal{L}_{\text{eff}}}{z_o^3} \right) \left[|\vec{m}_{\text{eff}}|^2 - \frac{1}{4 \pi^2 e^2 N_s^2} \left(\frac{\Delta\psi_e}{\theta} \right)^2 \right] \quad (14)$$

The above expression is the desired equation for the atom desorption energy.

Equation (14) is compared to the data of Taylor and Langmuir² on the variation in cesium atom desorption energy from a tungsten surface. The values for $|\vec{m}_{\text{eff}}|$ and \mathcal{L}_{eff} are determined from their work function data, and z_o is chosen so that the calculated values of $-\Delta\psi_a$ fit the data at medium and at high coverages. The various values used are

$$\begin{aligned} |\vec{m}_{\text{eff}}| &= 9.70 \text{ Debyes} \\ \mathcal{L}_{\text{eff}} &= 18.7 \text{ \AA}^3 \\ z_o &= 3.22 \text{ \AA} \\ N_s &= 4.8 \times 10^{14} / \text{cm}^2 \end{aligned} \quad (15)$$

Figure 3 shows the comparison of the calculated and experimental values of $-\Delta\psi_a$. Points for the solid line are computed from equation (14), and points for the dotted line are computed from the equation

$$\Delta\psi_a = - \frac{1}{4 \mathcal{L}_{\text{eff}}} \left[|\vec{m}_{\text{eff}}|^2 - \frac{1}{4 \pi^2 e^2 N_s^2} \left(\frac{\Delta\psi_e}{\theta} \right)^2 \right] \quad (16)$$

which is equation (14) with the term $\mathcal{L}_{\text{eff}}/z_o^3$ set equal to zero. Equation (16) corresponds to the case in which an adsorbed particle's own image field is not taken into account in the determination of the total dipole field acting on it.

It is interesting to note that the polarizability α_c of adsorbed cesium computed from equation (8) with the values of α_{eff} and z_o given in equation (15) is 6.00 \AA^3 , which is quite close to the ionic polarizability of cesium. Also, the value of $z_o/2$ is close to the ionic radius of cesium.

In this paper it has been shown that the correct polarizability which should be used in equation (7) is the effective polarizability defined by equation (8). In addition, it is demonstrated that the variation in atom desorption energy is closely related to the variation in electron work function with coverage.

e	electronic charge
\vec{E}	dipole field
\vec{E}_{eff}	effective dipole field
\vec{m}_{eff}	effective dipole moment as coverage approaches zero
\hat{n}	unit vector normal to and coming out of surface
N_s	number of adsorption sites per unit area
\vec{p}_{eff}	effective dipole moment at coverages greater than zero
T_{cs}	cesium reservoir temperature
T_s	surface temperature
$W_{s-\infty}$	work required to take a particle from the surface to infinity
$\Delta W_{s-\infty}$	change in work required to take a particle from the surface to infinity
$\Delta W_{\infty-s}$	change in work required to take a particle from infinity to the surface
z_0	distance between the center of an adsorbed particle and the center of its image
\mathcal{L}	polarizability of an adsorbed particle
\mathcal{L}_{eff}	effective polarizability of an adsorbed particle
θ	coverage, or the fraction of surface occupied by adsorbed particles
ϕ_a	atom desorption energy
$\Delta\phi_a$	change in atom desorption energy
$\Delta\phi_e$	change in electron work function


SUBSCRIPTS


i, k i th and k th adsorbed particles

REFERENCES

1. Houston, John M.: Thermionic Emission of Refractory Metals in Cesium Vapor. Power Information Center-Proceedings of the Round Table Discussion on Cathode Emission Investigations and Experimental Techniques For Fabrication and Operating Thermionic Cells. University of Pennsylvania, 1 and 2, June 1961. PIC-ELE-TI 3/3. June 30, 1961.
2. Taylor, John Bradshaw, and Langmuir, Irving: The Evaporation of Atoms, Ions, and Electrons from Cesium Films on Tungsten. Physical Review, vol. 44, no. 6, Sept. 15, 1933, pp. 423-458.

WORK FUNCTION VARIATION AS FUNCTION OF
RATIO OF SURFACE TO CESIUM TEMPERATURE





VARIATION IN THE Cs ATOM DESORPTION ENERGY FROM
A W SURFACE AS FUNCTION OF Cs COVERAGE

